

عدد المسائل : أربع	مسابقة في مادة الرياضيات المدة : ساعتان	الاسم: الرقم:
--------------------	--	------------------

ملاحظة : يُسمح باستعمال آلة حاسبة غير قابلة للبرمجة أو اختران المعلومات أو رسم البيانات.
يستطيع المُرشح الاجابة بالترتيب الذي يناسبه (دون الالتزام بترتيب المسائل الوارد في المسابقة).

I- (4 points)

An insurance company offers its employees a special fee for an annual life insurance policy. The following table represents this offer:

Age of the employee in years	25	26	27	28	29
Rank x_i	0	1	2	3	4
Annual fee y_i in hundred thousand LL	2	2.5	3.25	3.75	4

- 1) a- Calculate $\bar{x}$ and $\bar{y}$, the respective means (averages) of the two variables x and y .
b- Write an equation of (D), the regression line of y in terms of x .
- 2) Draw, in a rectangular system, the scatter plot of the points associated to the distribution $(x_i; y_i)$ as well as the center of gravity G and the line (D).
(Unit on the x -axis = 2 cm , unit on the y -axis = 4 cm.).
- 3) Assume that the above pattern remains valid till the age of 35. Estimate the annual fee for an employee whose age is 31 years.
- 4) What is the percentage of increase in the insurance annual fee between a 25 year old employee and a 31 year old employee?

II- (4 points)

The employees of a school are distributed into three categories: Instructors, administrators and technicians.

- 80% of the employees are instructors of which 70% are women.
- 10% of the employees are administrators of which 80% are women.
- 65% of the employees are women.

One employee is randomly selected from the school. Consider the following events:

- I: « The employee selected is an instructor »
A: « The employee selected is an administrator »
T: « The employee selected is a technician »
M: « The employee selected is a man »
W: « The employee selected is a woman ».

- 1) Calculate the probability $P(M)$.
- 2) Calculate $P(I \cap W)$; $P(A \cap W)$ and verify that $P(W / T) = 0.1$.
- 3) Knowing that the selected employee is a man, what is the probability that he is an instructor?
- 4) In this part, suppose that the number of employees in this school is 200. The names of these employees are written on cards and these cards are placed in a box.

Three cards are randomly and simultaneously selected from this box.

What is the probability of the event B: « None of the three cards holds the name of a technician, at least one card holds the name of an administrator and at least one card holds the name of an instructor »?

III- (4 points)

At the end of July 2015, a retiree deposits an amount of 220 million LL in a bank at an annual interest rate of 6% compounded monthly. At the end of each month and after the compounding of interest, this person decides to withdraw 2 million LL for his expenses.

Denote by the natural number n the rank of the month after the deposit and by U_n the amount, in millions LL, that this person has in his account in the n th month after the withdrawal of the 2 million LL.

Thus $U_0 = 220$.

- 1) Verify that $U_1 = 219.1$ and calculate U_2 .
- 2) For all natural numbers n , show that $U_{n+1} = 1.005U_n - 2$.
- 3) For all natural numbers n , let $V_n = U_n - 400$.
 - a- Prove that (V_n) is a geometric sequence. Specify its ratio and first term.
 - b- Show that $U_n = -180 \times (1.005)^n + 400$.
 - c- Prove that the sequence (U_n) is decreasing.
 - d- In how many years would the amount in the bank account become less than half the original amount for the first time?

IV- (8 points)

Part A

Consider the function f defined over $[0; +\infty[$ as $f(x) = -x^2 e^{-x} + 3$ and denote by (C) its representative curve in an orthonormal system $(O; \vec{i}, \vec{j})$.

- 1) Determine $\lim_{x \rightarrow +\infty} f(x)$. Deduce an asymptote (d) to (C).
- 2) Show that $f'(x) = x(x-2)e^{-x}$ and set up the table of variations of the function f .
- 3) Draw (d) and (C).
- 4) Let F be the function defined over $[0; +\infty[$ as $F(x) = (x^2 + 2x + 2)e^{-x} + 3x$.
 - a- Prove that F is an antiderivative of the function f .
 - b- Deduce the area of the region bounded by the curve (C), the x -axis, the y -axis and the line with equation $x = 2$.

Part B

A factory produces paint. All the production is sold. The average cost of production in hundreds of thousands LL is given as $f(x) = -x^2 e^{-x} + 3$, where x is expressed in hundreds of liters. $x \in [0.2; 9]$.

- 1) Determine, in LL, the average cost for the production of 600 liters of paint.
- 2) a- How many liters of paint should the factory produce so that the average cost of production is minimal? What is then the average cost in LL?
 - b- If the sale price of 100 liters of paint is 230 000 LL, does the factory make profit? Explain.
- 3) a- Express the total cost $C_T(x)$ in terms of x .
 - b- How many liters of paint should the factory produce so that the average cost of production is equal to the marginal cost?

I- (7 points)

Q	Correction	Grade
1-a	$\bar{x} = 2$; $\bar{y} = 3.1$	1
1-b	(D): $y = 0.525x + 2.05$	1
2	G(2,3,1)	2
3	age of 31 $\rightarrow x = 6$ then $y = 0.525(6) + 2.05 = 5.2$ therefore 520 000 LL	1.5
4	$\frac{5.2 - 2}{2} = 1.6$ then 160% is the increasing percentage	1.5

II- (7 points)

Q	Correction	Grade
1	$P(M) = 1 - P(W) = 1 - 0.65 = 0.35$	1
2	$P(I \cap W) = P\left(\frac{W}{I}\right) \times P(I) = 0.7 \times 0.8 = 0.56$; $P(A \cap W) = P\left(\frac{W}{A}\right) \times P(A) = 0.8 \times 0.1 = 0.08$; $P(W \cap T) = P(W) - P(W \cap I) - P(W \cap A) = 0.65 - 0.56 - 0.08 = 0.01$ $P\left(\frac{W}{T}\right) = \frac{P(W \cap T)}{P(T)} = \frac{0.01}{0.1} = 0.1$.	3
3	$P\left(\frac{I}{M}\right) = \frac{P(I \cap M)}{P(M)} = \frac{0.8 \times 0.3}{0.35} = 0.68$.	1.5
4	$P(B) = \frac{C_{160}^2 \times C_{20}^1 + C_{160}^1 \times C_{20}^2}{C_{200}^3} = \dots = \frac{1424}{6567} \approx 0.216$.	1.5

III- (7 points)

Q	Correction	Grade
1	$U_1 = \left(1 + \frac{0,06}{12}\right) \times 200 = 219,1$ et $U_2 = \left(1 + \frac{0,06}{12}\right) \times 219,1 = 218.1955$.	1
2	$U_{n+1} = \left(1 + \frac{0,06}{12}\right) \times U_n - 2 = 1.005 U_n - 2$	1
3-a	$V_{n+1} = U_{n+1} - 400 = 1.005 U_n - 402 = 1.005 (U_n - 400) = 1.005 V_n$ $q = 1.005$ and $V_0 = U_0 - 400 = 220 - 400 = -180$	1.5
3-b	$V_n = V_0 \times q^n = -180 \times (1.005)^n$ and $U_n = V_n + 400 = -180 \times (1.005)^n + 400$	1
3-c	$U_{n+1} - U_n = -180 \times (1.005)^{n+1} + 400 + 180 \times (1.005)^n - 400$ $= -180 \times (1.005)^n (1.005 - 1) = -0.9 \times (1.005)^n < 0$	1
3-d	$U_n < \frac{U_0}{2}$; $-180 \times (1.005)^n + 400 < 110$; $(1.005)^n > \frac{29}{18}$; $n > \frac{\ln(29/18)}{\ln(1.005)}$; $n > 95.6$; $n = 96$ then after 8 years.	1.5

IV- (14 points)

Q	Correction	Grade														
A-1	$\lim_{x \rightarrow +\infty} f(x) = 3$ then (d) : $y = 3$ is a horizontal asymptote at $+\infty$	1														
A-2	$f'(x) = x(x-2)e^{-x}$ same sign as $x-2$ then: <table><tr><td>x</td><td>0</td><td>2</td><td>$+\infty$</td></tr><tr><td>$f'(x)$</td><td>0</td><td>—</td><td>0</td><td>+</td></tr><tr><td>$f(x)$</td><td>3</td><td></td><td>$3 - \frac{4}{e^2}$</td><td>3</td></tr></table>	x	0	2	$+\infty$	$f'(x)$	0	—	0	+	$f(x)$	3		$3 - \frac{4}{e^2}$	3	1.5
x	0	2	$+\infty$													
$f'(x)$	0	—	0	+												
$f(x)$	3		$3 - \frac{4}{e^2}$	3												
A-3		1.5														
A-4-a	$F'(x) = (2x+2)e^{-x} - e^{-x}(x^2+2x+2) + 3 = f(x)$	1.5														
A-4-b	$A = \int_0^2 f(x) dx = [F(x)]_0^2 = F(2) - F(0) = \frac{10}{e^2} + 6 - 2 = \left(4 + \frac{10}{e^2}\right)u^2$	1.5														
B-1	$f(6) = -36e^{-6} + 3 = 2.910765$ the average cost is: 2 910 765 LL.	1.5														
B-2-a	According to the table of variations the average cost production is minimal for $x = 2$ then 200 liters. The minimal average cost is $f(2) = 2.45866$ then 245 866 LL	1.5														
B-2-b	$y = 2,3 < f(2)$ (The minimum of f) then no gain.	1.5														
B-3-a	$C_T(x) = x f(x) = -x^3 e^{-x} + 3x$.	1														
B-3-b	$C_m(x) = (x^3 - 3x^2)e^{-x} + 3$; $C_M(x) = C_m(x)$; $(x^3 - 3x^2)e^{-x} + 3 = -x^2 e^{-x} + 3$; $x^3 - 3x^2 = -x^2$; $x^2(x-2) = 0$; $x = 0$ or $x = 2$ but $0 \notin [0.2 ; 9]$ then $x = 2$ therefore for a production of 200 liters.	1.5														

||